



Sharpes Pullet Grower Pellets

SHARPES PULLET GROWER PELLETS ARE MADE WITH 100% NZ GRAINS. This is a complete ration specifically formulated for young layer pullets not in lay.

KEY BENEFITS:

- This ration contains a coccidiostat to aid in the prevention of coccidiosis in young birds.
- Formulated with high quality, locally sourced ingredients, with the addition of meat and bone meal for an excellent source of protein for growing birds.

BAG SIZES: 10kg, 20kg, 1/2 tonne and 1 tonne bags.

FEED RECOMMENDATION: Layer Pullets - Feed ad lib from 6 to 8 weeks of age until the point of lay, typically around 17 weeks.

TOP TIP: Always ensure that birds have access to fresh, clean water.

INGREDIENTS SELECTED FROM: Locally sourced grains and cereals; meat and bone meal, plant proteins, limestone, macro and trace minerals δ vitamins, RumensinTM 20% MillMixTM.

TYPICAL ANALYSIS (approximate on an as fed basis):

Crude Protein 19% Fat 3.0%

STORAGE: To maintain quality please store product in a cool, dry and vermin free environment. Best fed within six months of production.

WARNING This feed contains Rumensin™ 20% Millmix™.

For Animal Treatment Only. Do not feed to Sheep, Cattle, Deer, Goats, Buffaloes, or other ruminant animals. This product contains or may contain Ruminant Protein.

This feed contains Rumensin™ 20% Millmix™ in the poultry feed at a rate of 0.5 kg premix per tonne of feed as an aid in the prevention of coccidiosis caused by Eimeria acervulina, E.brunetti, E.maxima, E.mivati, E.necatrix and E.tenella. This is the equivalent of 100 ppm (mg/kg) of monensin active ingredient in the poultry feed.

Rumensin[™] 20% MillMix[™] is a registered pursuant to the ACVM Act 1997, A009107. See www.nzfsa.govt.nz/acvm for registration conditions.

By law the user must take due care, obtaining expert advice, when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.







Sharpes Pullet Grower Pellets with Coccidiostat

CONTRAINDICATIONS: Poultry consuming Rumensin™ 20% MillMix™ should not be treated with products containing tiamulin or oleandomycin. Severe growth depression may occur.

CAUTION: Do not use on horses, dogs or other equid or camelid species as fatal toxicosis may result. Ensure recommended doses are not exceeded. Care must be exercised when feeding concurrently with other antimicrobials.

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

WITHHOLDING PERIOD:

Poultry: Meat – Nil Eggs – 10 days

Please note: Do not feed to any other animal species than those listed on the label and only feed as directed.

Nutritional information provided by: Nutritech International.







